

Supplementary Materials

Anti-infection effects of heparin on SARS-CoV-2 in a diabetic mouse model

Zhongyun Zhang^{1,2,#}, Ning Zhang^{3,#}, Xuancheng Lu^{4,#}, Min Zhou⁵, Xiaoxiang Yan⁶,
Weiqiong Gu^{1,2}, Jingru Yang⁷, Qin Zhang⁴, Cheng Zhang³, Yuhuan Gong³, Mingjun
Jia³, Xiaoyu Zhang³, Peng Ning³, Mei Liu⁴, Xiaoyan Li⁴, Xiaomeng Shi⁴, Wenjun Liu^{3,8},
George F. Gao^{3,4,8}, Guang Ning^{1,2}, Jiqiu Wang^{1,2,*}, Yuhai Bi^{3,7,8,*}

¹Department of Endocrinology and Metabolism, China National Research Center for Metabolic Diseases, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

²Shanghai National Clinical Research Center for Metabolic Diseases, Key Laboratory for Endocrine and Metabolic Diseases of the National Health Commission of the PR China, Shanghai National Center for Translational Medicine, Shanghai 200025, China

³CAS Key Laboratory of Pathogen Microbiology and Immunology, Institute of Microbiology, Center for Influenza Research and Early-warning (CASCIRE), CAS-TWAS Center of Excellence for Emerging Infectious Diseases (CEEID), Chinese Academy of Sciences, Beijing 100101, China

⁴Laboratory Animal Center, Chinese Center for Disease Control and Prevention (China CDC), Beijing 102206, China

⁵Department of Respiratory and Critical Care Medicine, Institute of Respiratory Diseases, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200025, China

⁶Department of Cardiology, Institute of Cardiovascular Diseases, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200025, China

⁷School of Laboratory Medicine and Life Sciences, Wenzhou Medical University, Wenzhou, Zhejiang 325000, China

⁸University of Chinese Academy of Sciences, Beijing 100049, China

#Authors contributed equally to this work

*Corresponding authors, E-mail: wangjq@shsmu.edu.cn; beeyh@im.ac.cn

Supplementary Table S1. Characteristics of COVID-19 patients with vs without diabetes at baseline.

Characteristic	Non-diabetes (291)		Diabetes (66)		t/Z/x ²	Test	P
	Mean ± SD or median (IQR) or n (%)	Missing Data	Mean ± SD or median (IQR) or n (%)	Missing Data			
Age, year, median (IQR)	64.00 (49.00-70.00)	0	68.00 (59.75–73.25)	0	-2.933	Mann-Whitney test	0.003
Sex		0		0	1.120	Chi-Square tests	0.290
Male, n (%)	151 (51.9%)	-	39 (59.1%)	-			
Female, n (%)	140 (48.1%)	-	27 (40.9%)	-			
Smoking, n (%)	16 (7.1%)	65	5 (10.2%)	17	-	Fisher's exact test	0.551
Comorbidities							
Hypertension, n (%)	92 (31.6%)	0	39 (59.1%)	0	17.483	Chi-Square tests	<0.001
Hyperlipemia, n (%)	3 (1.0%)	0	3 (4.5%)	0	-	Fisher's exact test	0.079
CHD, n (%)	27 (9.3%)	0	14 (21.2%)	0	7.537	Chi-square tests	0.006
COPD, n (%)	18 (6.2%)	0	8 (12.1%)	0	-	Fisher's exact test	0.113
CKD, n (%)	7 (2.4%)	0	5 (7.6%)	0	-	Fisher's exact test	0.051
Cancer, n (%)	10 (3.4%)	0	2 (3.0%)	0	-	Fisher's exact test	1.000
Main symptoms of COVID-19							
Cough, n (%)	234 (80.4%)	0	56 (84.8%)	0	0.694	Chi-square tests	0.405
Fever, n (%)	225 (77.3%)	0	50 (75.8%)	0	0.074	Chi-square tests	0.785
Expectoration, n (%)	153 (52.6%)	0	44 (66.7%)	0	4.318	Chi-square tests	0.038
Main laboratory findings							
FBG, mmol/L, median (IQR)	5.45 (4.96–6.67)	27	7.35 (5.89–9.85)	7	-5.659	Mann-Whitney test	<0.001
Hemoglobin, g/L, median (IQR)	128.00 (116.00–137.00)	3	126.00 (115.00–139.00)	3	-0.627	Mann-Whitney test	0.531
RBC count, 10 ⁹ /L, mean ± SD	4.09 ± 0.61	3	4.08 ± 0.56	3	0.046	t-test	0.964

WBC count, 10 ⁹ /L, median (IQR)	5.73 (4.45–7.37)	4	6.14 (4.97–7.75)	3	-1.452	Mann-Whitney test	0.146
Lymphocyte count, 10 ⁹ /L, mean ± SD	1.27 ± 0.55	3	1.29 ± 0.60	3	-0.217	<i>t</i> -test	0.828
Neutrophil count, 10 ⁹ /L, median (IQR)	3.68 (2.59–5.20)	3	4.02 (2.86–5.87)	3	-1.469	Mann-Whitney test	0.142
Monocyte count, 10 ⁹ /L, median (IQR)	0.47 (0.36–0.63)	3	0.49 (0.37–0.67)	3	-1.069	Mann-Whitney test	0.285
Platelet count, 10 ⁹ /L, mean ± SD	248.55 ± 104.45	3	251.39 ± 91.06	4	-0.199	<i>t</i> -test	0.843
CRP, mg/L, median (IQR)	12.00 (1.63–59.10)	11	8.60 (2.20–80.65)	1	-0.449	Mann-Whitney test	0.654
Signs at admission							
SpO ₂ , %, median (IQR)	97.00 (95.00–99.00)	27	96.00 (92.00–98.00)	7	-2.366	Mann-Whitney test	0.018
RR, bpm, median (IQR)	20.00 (18.00–22.00)	15	20.00 (20.00–22.00)	3	-1.976	Mann-Whitney test	0.048
HR, bpm, median (IQR)	86.50 (78.00–96.00)	11	89.50 (76.00–102.00)	2	-0.660	Mann-Whitney test	0.509
BT, °C, median (IQR)	36.70 (36.30–37.00)	19	36.60 (36.30–37.00)	3	-0.923	Mann-Whitney test	0.356
Death, n (%)	47 (16.2%)	0	9 (13.6%)	0	0.257	Chi-square tests	0.612

Note: Bold values represent statistically significant P values.

Abbreviations: CHD, coronary heart disease; COPD, chronic obstructive pulmonary disease; CKD, chronic kidney disease; FBG, fasting blood glucose; RBC, red blood cell; WBC, white blood cell; CRP, C-reactive protein; SpO₂; oxygen saturation; RR, respiratory rate; HR, heart rate; BT, body temperature; IQR, interquartile range; SD, standard deviation.

Supplementary Table S2. Characteristics of COVID-19 patients with vs without diabetes at baseline (other clinical features with no statistically significant difference).

Characteristic	Non diabetes (389)		Diabetes (86)		t/Z/x ²	Test	P
	Mean ± SD or median (IQR) or n (%)	Missing data	Mean ± SD or median (IQR) or n (%)	Missing Data			
Main symptoms of COVID-19							
Hemoptysis, n (%)	8 (2.7%)	0	0 (0.0%)	0	-	Fisher's exact test	0.360
Shortness of breath, n (%)	106 (36.4%)	0	23 (34.8%)	0	0.058	Chi-square tests	0.810
Pharyngalgia, n (%)	37 (12.7%)	0	9 (13.6%)	0	0.041	Chi-square tests	0.840
Headache, n (%)	43 (14.8%)	0	10 (15.2%)	0	0.006	Chi-square tests	0.938
Dizzy, n (%)	19 (6.5%)	0	3 (4.5%)	0	-	Fisher's exact test	0.777
Muscular soreness, n (%)	62 (21.3%)	0	14 (21.2%)	0	<0.001	Chi-square tests	0.987
Weakness, n (%)	117 (40.2%)	0	24 (36.4%)	0	0.332	Chi-square tests	0.564
Pectoralgia, n (%)	56 (19.2%)	0	10 (15.2%)	0	0.598	Chi-square tests	0.439
Chest stuffiness, n (%)	126 (43.3%)	0	19 (28.8%)	0	4.697	Chi-square tests	0.030
Diarrhea, n (%)	71 (24.4%)	0	15 (22.7%)	0	0.082	Chi-square tests	0.774
Nausea and vomiting, n (%)	41 (14.1%)	0	11 (16.7%)	0	0.287	Chi-square tests	0.592
Palpitation, n (%)	6 (2.1%)	0	2 (3.0%)	0	-	Fisher's exact test	0.644
Main laboratory findings							
PT, s, median (IQR)	13.70 (13.20–14.40)	17	13.75 (13.00–14.50)	6	-0.197	Mann-Whitney test	0.844
PTA, s, median (IQR)	91.50 (82.25–98.00)	19	90.00 (81.00–100.75)	6	-0.166	Mann-Whitney test	0.868
INR, median (IQR)	1.06 (1.01–1.14)	18	1.07 (0.99–1.14)	6	-0.051	Mann-Whitney test	0.959
Fg, g/L, median (IQR)	4.25 (3.40–5.36)	18	4.33 (3.62–5.75)	6	-0.82	Mann-Whitney test	0.412
APTT, s, median (IQR)	37.80 (35.25–41.10)	18	38.40 (36.00–41.05)	6	-0.742	Mann-Whitney test	0.458
TT, s, median (IQR)	16.60 (15.75–17.50)	18	16.70 (15.83–17.70)	6	-0.285	Mann-Whitney test	0.776

D dimer, median (IQR)	0.79 (0.32–2.47)	23	0.58 (0.36–2.95)	10	-0.296	Mann-Whitney test	0.767
FDP, µg/mL, median (IQR)	4.00 (4.00–9.20)	34	4.00 (4.00–9.875)	14	-0.180	Mann-Whitney test	0.857
ATIII, %, median (IQR)	92.00 (81.00–101.00)	37	91.00 (85.00–98.00)	14	-0.355	Mann-Whitney test	0.722
Complications							
RF, n (%)	23 (7.9%)	0	3 (4.5%)	0	-	Fisher's exact test	0.439
ARDS, n (%)	41 (14.1%)	0	10 (15.2%)	0	0.05	Chi-square tests	0.824
AMI, n (%)	54 (18.6%)	0	8 (12.1%)	0	1.553	Chi-Square tests	0.213
ALI, n (%)	63 (21.6%)	0	12 (18.2%)	0	0.39	Chi-Square tests	0.532
AKI, n (%)	29 (10.0%)	0	10 (15.2%)	0	1.487	Chi-Square tests	0.223
Septic shock, n (%)	17 (5.8%)	0	7 (10.6%)	0	-	Fisher's exact test	0.175
MODS, n (%)	3 (1.0%)	0	0 (0.0%)	0	-	Fisher's exact test	1.000

Note: Bold values represent statistically significant P values.


Abbreviations: PT, prothrombin time; PTA, prothrombin activity; INR, international normalized ratio; Fg, fibrinogen; APTT, activated Partial thromboplastin time; TT, thrombin time; FDP, fibrinogen degradation products; AT III, antithrombin III; RF, respiratory failure; ARDS, acute respiratory distress syndrome; AMI, acute myocardial injury; ALI, acute liver injury; AKI, acute kidney injury; MODS, multiple organ dysfunction syndrome; IQR, interquartile range.

Supplementary Table S3. Univariate and multivariate logistic regression analysis between fasting blood glucose and incidence of main symptoms and important complications in COVID-19 patients.

Characteristic	Crude OR (95% CI)	P	Adjust OR (95% CI)	P
Main symptoms of COVID-19				
Cough	1.04 (0.94–1.15)	0.427	-	-
Fever	1.02 (0.94–1.12)	0.584	-	-
Expectoration	1.04 (0.97–1.12)	0.256	-	-
Hemoptysis	1.03 (0.85–1.26)	0.757	-	-
Shortness of breath	1.01 (0.95–1.09)	0.678	-	-
Pharyngalgia	1.08 (1.00–1.16)	0.051	-	-
Headache	0.99 (0.90–1.09)	0.853	-	-
Dizzy	1.03 (0.92–1.16)	0.574	-	-
Muscular soreness	1.00 (0.92–1.08)	0.928	-	-
Weakness	1.02 (0.96–1.09)	0.502	-	-
Pectoralgia	1.03 (0.95–1.11)	0.531	-	-
Chest stuffiness	1.03 (0.96–1.10)	0.406	-	-
Diarrhea	1.00 (0.92–1.08)	0.955	-	-
Nausea and vomiting	1.02 (0.94–1.11)	0.624	-	-
Palpitation	1.03 (0.86–1.24)	0.724	-	-
Complications				
RF	1.11 (1.03–1.21)	0.012	1.10 (1.00–1.20)	0.041
ARDS	1.15 (1.06–1.24)	<0.001	1.12 (1.04–1.21)	0.003
AMI	1.10 (1.02–1.18)	0.009	-	-
ALI	1.03 (0.96–1.11)	0.430	-	-
AKI	1.14 (1.06–1.24)	0.001	1.11 (1.03–1.20)	0.006
Septic shock	1.11 (1.02–1.20)	0.016	1.09 (1.00–1.19)	0.055
MODS	0.87 (0.39–1.97)	0.741	-	-
Death	1.18 (1.09–1.29)	0.001	1.15 (1.06–1.24)	0.001


Note: Bold values represent significant P values.

Abbreviations: RF, respiratory failure; ARDS, acute respiratory distress syndrome; AMI, acute myocardial injury; ALI, acute liver injury; AKI, acute kidney injury; MODS, multiple organ dysfunction syndrome; OR, odds ratio.



Supplementary Figure S1 hACE2-transgenic mice are sensitive to SARS-CoV-2 infection under a chow diet (CD) and high-fat diet (HFD)

F4/80 in lungs of hACE2-CD and hACE2-HFD mice after infection was detected using immunohistochemistry (IHC). 5×, scale bar represents 300 μm, 20×, scale bar represents 100 μm.



Supplementary Figure S2 Diabetic mice exhibit more severe symptoms after SARS-CoV-2 infection

F4/80 in lungs of hACE2-Lepr^{-/-} and hACE2-Lepr^{+/+} mice after infection was detected using IHC. 5×, scale bar represents 300 μm, 20×, scale bar represents 100 μm.