

Supplementary Materials

Supplementary Materials and Methods

Sampling

During 2014 and 2017, we conducted a joint Sino-Kazh field investigation on the herpetofauna in eastern Kazakhstan and the adjacent Junggar Depression in China. Thirty specimens (15 males, 15 females) from seven sites were collected from eastern Kazakhstan and northwestern Altay region, Xinjiang, China (Supplementary Figure S1 and Table S1), which were initially identified as *E. multiocellata*. The lizards were euthanized with an overdose of sodium pentobarbital via intraperitoneal injection, and liver or tail tissues were extracted and preserved in 95% ethanol following the animal use protocols approved by the Chengdu Institute of Biology (CIB), Chinese Academy of Sciences (CAS). All voucher specimens are preserved in CIB, CAS.

Molecular data collection

Genomic DNA was extracted from liver or tail tissues following the modified high-salt protocol proposed by Aljanabi & Martinez (1997). Fragments of the mitochondrial *COI* gene (709 bp) were amplified via polymerase chain reaction (PCR) using primers designed according to Ward et al. (2005) with minor modifications (E_COIF 5'-TCAACCAACCACAAAGACATTGGCAC-3' and E_COIR 5'-TAGACTTCTGGGTGGCCAAAGAATCA-3'). The PCR reaction mixture contained 12.5 µL of 2 × EasyTaq SuperMix (Tsingke Biol-Tech, Chengdu, China), 0.2 µM of each primer, and 1–2 µL of genomic DNA for a total volume of 25 µL. The amplification protocols included an initial denaturation at 94 °C for 4 min, followed by 35 cycles of 94 °C for 30 s, 52 °C for 30 s and elongation at 72 °C for 50 s, and a final extension at 72 °C for 10 min. All PCR products were commercially purified and sequenced for double strands with the primers used for amplification. All novel sequences were deposited in GenBank under the accession numbers MW172523–MW172552.

Nineteen individuals from China and 11 from Kazakhstan were sequenced for *COI*. All novel sequences were translated to amino acids with MEGA v7.0.26 (Kumar et al., 2016), and no stop codons were detected. In total, 156 sequences – which included those from Orlova et al. (2017) (available in GenBank) – were aligned using Clustal X v2.0 (Larkin et al., 2007) with default settings. Identical sequences were collapsed into a single haplotype using DnaSP v5.0 (Librado & Rozas, 2009).

DNA barcoding diversity analysis

The dataset containing all sequences was uploaded into the BOLD system under the project ‘*Eremias multiocellata-przewalskii* complex barcoding’ (Barcode of Life Data System website, www.barcodinglife.com; Ratnasingham & Hebert, 2007). We measured the genetic distance with the Kimura 2-parameter (K2P; Kimura, 1980) model, which has been widely used in barcoding studies. The intraspecific and intra-genus pairwise distances were calculated with the ‘Distance Summary’ analysis tool in BOLD. The maximum intraspecific genetic distance and corresponding minimum distance to nearest neighbor at the species level were calculated and compared using the ‘Barcoding Gap Analysis’ tool in BOLD.

Phylogenetic analysis

The neighbor-joining (NJ) (Saitou & Nei, 1987) tree was constructed in MEGA with a gamma distribution rate variation among sites (shape parameter = 4) and pairwise deletion option for the treatment of missing data. Bootstrap support for clades was evaluated by nonparametric bootstrap analysis (Felsenstein, 1985) with 1 000 replicates. Bayesian inference (BI) was conducted with MrBayes v3.2.2 (Ronquist et al., 2012). The best-fit models of nucleotide substitution for each partition scheme were selected using PartitionFinder v2.1.1 (Lanfear et al., 2017). Three codon partitions and their corresponding substitution model for the *COI* gene sequences were proposed: first codon, K80+G+I; second codon, F81; third codon, GTR+G. Two parallel runs of one cold and three heated Monte Carlo Markov chains (MCMCs) were performed with sampling every 200 generations and were automatically stopped

after reaching a mean standard deviation between split frequencies of less than 0.01. Parameters and topologies were estimated from 13 405 steps after discarding the first 30%. Convergence of the runs was assessed by effective sample size ($\text{ESS} \geq 200$) with Tracer v1.7.1 (Rambaut et al., 2018). Partitioned ML (Maximum Likelihood) analyses were conducted in RAxML v8.2.4 (Stamatakis, 2014) with the GTR+G model for all subsets. A complete random starting tree (option -d) was initiated with 100 independent search replicates where the tree with the best likelihood was chosen. Bootstrap support proportion (BSP) for the clades was obtained with a sufficient number of bootstrap replicates (450), which was automatically determined by setting the default cutoff threshold (option -# autoMRE).

Species delimitation

We first employed two distance-based methods dedicated to barcoding datasets. The Barcode Index Number (BIN) system clusters (Ratnasingham & Hebert, 2013) was conducted with the ‘BIN Discordance’ option in BOLD, which was employed to assign individuals to operational taxonomic units (OTUs) that represent presumable species. The BIN system uses the refined single linkage algorithm (RESL) for clustering process, and hence is independent of any prior taxonomic assignment. Similarly, Automatic Barcode Gap Discovery (ABGD; Puillandre et al., 2012) does not need *a priori* species hypotheses, and clusters sequences into candidate species by detecting a ‘barcode gap’ that demarcates intra- and interspecific distance. Analysis was implemented in the ABGD web server (<https://bioinfo.mnhn.fr/abi/public/abgd/abgdweb.html>) with the K2P model and other parameters as default.

Two coalescent methods were further employed to delimit species: the generalized mixed Yule coalescent (GMYC) model implemented in the GMYC web server (<http://species.h-its.org/gmyc/>), and Bayesian species identification under the multispecies coalescent (MSC) model with the program BPP v3.3 (Yang & Rannala, 2010). GMYC was proposed to delimit species using the reconstructed gene tree for a single locus (Fujisawa & Barraclough, 2013). We reconstructed ultrametric gene trees

in BEAST v1.8.2 (Drummond et al., 2012) with three codon partitions. The substitution models were calculated in PartitionFinder; the first codon used K80+G+I, the second used HKY, and the third used TRN+G. To reduce compounding errors from time calibration uncertainties or molecular substitution rates, we left the prior ‘clock.rate’ to default 1.0. Posterior samplings were drawn every 1 000 steps over 15 million MCMC steps with the uncorrelated lognormal relaxed clock model and Yule speciation tree prior. Two independent runs were performed. Stationarity of the Markov chains for each run was checked using Tracer for $\text{ESS} \geq 200$. Finally, the tree files were combined using LogCombiner v1.8.2, and a maximum clade credibility (MCC) tree with median node heights was generated with TreeAnnotator after discarding the first 25% of posterior sampling as burn-in. The MCC tree was used as an input in the single-threshold GMYC analysis implemented in the GMYC server. Bayesian species identification under the MSC model was considered to have better performance in species assignment than other single-locus delimiting methods for DNA barcoding analyses (Yang & Rannala, 2017). We ran the A11 model with the presumption of relatively large ancestral population sizes ($\theta \sim \text{G}(1, 200)$) and deep divergence ($\tau_0 \sim \text{G}(1.5, 10)$); this single set of priors was based on our unpublished data (Liu, 2019). To explore the potential cryptic species within our study system, we designated the significantly supported lineages within *E. stummeri* (two reciprocal monophyletic lineages) and *Eremias* sp. 3 (two reciprocal monophyletic lineages and a singleton) as different unique populations in our prior population assignments in the imap file. The reversible-jump Markov chain Monte Carlo (rjMCMC) algorithm 1 with parameters combination (α, m) = (1, 0.5) and speciesmodelprior = 1 were used to repeat the BPP runs twice with different starting seeds to assess the consistency between the different runs. Each run was conducted with rjMCMC analysis for 500 000 generations (sampling interval of five), with the first 20 000 samples discarded as burn-in.

In addition, species delimitation with the multi-rate Poisson tree processes model (mPTP) has been shown to have reliable performance (Blair & Bryson, 2017), as it explicitly accounts for differences in sampling intensity and/or effective population

sizes of species (Kapli et al., 2017). We implemented mPTP analysis in the web server (<https://mptp.h-its.org>) using the MCC tree constructed in the previous BEAST analysis.

All species delineation analyses described above were conducted with haplotypes (except BIN, which used all sequences) and without outgroups.

Morphological analyses

Morphological measurement was performed on all 30 specimens collected in this study, following the traits analyzed in Szczerbak (1974), Dujsebayeva et al. (2009), and Orlova et al. (2017). Morphological analyses were conducted with 11 metric and 12 meristic traits (Supplementary Table S5). Of these traits, four meristic traits (Supplementary Table S5) were counted on both the left and right sides of the body. Metric measurements were conducted using a digital caliper and recorded to two decimal places.

We first employed analysis of variance (ANOVA) to test for sexual dimorphism in both metric and meristic datasets collected in this study. All metric traits (except Dist.P.fm) and one meristic trait (Sq.c.cd) exhibited significant sexual dimorphism ($p < 0.05$). As a result, the comparative demonstration of these sex-dependent characters (values given as ranges and means \pm SE) between the individuals from this study and the holotype and paratypes of *E. dzungarica* from Orlova et al. (2017) were given separately for males and females. The values of the other age- and sex-independent metric and meristic characters were calculated for all specimens combined (including males, females and juveniles in this study and Orlova et al. (2017), respectively). Moreover, to test whether the specimens in this study differed significantly from *E. dzungarica* in morphological traits with a sample size greater than 11, we performed a two independent samples *t*-test. All morphometric analyses were performed with SPSS 19 (SPSS Inc., Chicago, USA).

REFERENCES


- Aljanabi SM, Martinez I. 1997. Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques. *Nucleic Acids Research*, **25**(22): 4692–4693.
- Blair C, Bryson Jr RW. 2017. Cryptic diversity and discordance in single-locus species delimitation methods within horned lizards (Phrynosomatidae: *Phrynosoma*). *Molecular Ecology Resources*, **17**(6): 1168–1182.
- Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution*, **29**(8): 1969–1973.
- Dujsebayeva TN, Chirikova MA, Belyalov OV. 2009. New finds of the racerunner of *Eremias multiocellata* complex in Kazakhstan. *Russian Journal of Herpetology*, **16**(1): 51–56.
- Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, **39**(4): 783–791.
- Fujisawa T, Barraclough TG. 2013. Delimiting species using single-locus data and the Generalized Mixed Yule Coalescent approach: a revised method and evaluation on simulated data sets. *Systematic Biology*, **62**(5): 707–724.
- Kapli P, Lutteropp S, Zhang J, Kobert K, Pavlidis P, Stamatakis A, et al. 2017. Multi-rate Poisson tree processes for single-locus species delimitation under maximum likelihood and Markov chain Monte Carlo. *Bioinformatics*, **33**(11): 1630–1638.
- Kimura M. 1980. A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, **16**(2): 111–120.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, **33**(7): 1870–1874.
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B. 2017. Partitionfinder 2:

- new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Molecular Biology and Evolution*, **34**(3): 772–773.
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGgettigan PA, McWilliam H, et al. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics*, **23**(21): 2947–2948.
- Librado P, Rozas J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, **25**(11): 1451–1452.
- Liu J. 2019. Molecular Phylogeny and Phylogeography of *Eremias velox* Complex and *Eremias multiocellata* Complex. Ph.D. dissertation, Sichuan University, Chengdu. (in Chinese)
- Orlova VF, Poyarkov Jr NA, Chirikova MA, Nazarov RA, Munkhbaatar M, Munkhbayar K, et al. 2017. MtDNA differentiation and taxonomy of Central Asian racerunners of *Eremias multiocellata*-*E. przewalskii* species complex (Squamata, Lacertidae). *Zootaxa*, **4282**(1): 1–42.
- Puillandre N, Lambert A, Brouillet S, Achaz G. 2012. ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology*, **21**(8): 1864–1877.
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018. Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology*, **67**(5): 901–904.
- Ratnasingham S, Hebert PDN. 2007. BOLD: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Molecular Ecology Notes*, **7**(3): 355–364.
- Ratnasingham S, Hebert PDN. 2013. A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. *PLoS One*, **8**(7): e66213.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, et al. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology*, **61**(3): 539–542.
- Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, **4**(4):

406–425.


- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, **30**(9): 1312–1313.
- Szczerbak NN. 1974. Yashchurki Palearktiki (*Eremias* Lizards of the Palearctic). Kiev: Naukova Dumka Press. (in Russian)
- Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PDN. 2005. DNA barcoding Australia's fish species. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **360**(1462): 1847–1857.
- Yang ZH, Rannala B. 2010. Bayesian species delimitation using multilocus sequence data. *Proceedings of the National Academy of Sciences of the United States of America*, **107**(20): 9264–9269.
- Yang ZH, Rannala B. 2017. Bayesian species identification under the multispecies coalescent provides significant improvements to DNA barcoding analyses. *Molecular Ecology*, **26**(11): 3028–3036.

Figure S1




Supplementary Figure S1 Frequency distribution of pairwise Kimura 2-parameter (K2P) distances (%) at intra- and interspecific taxonomic level in *Eremias multiocellata-przewalskii* species complex

Figure S2



Supplementary Figure S2 Scatterplot of maximum intraspecific variation and minimum genetic distance to the nearest-neighbor species for each species in *Eremias multiocellata-przewalskii* species complex


Fig. S3



Supplementary Figure S3 50% majority-rule consensus tree of *E. multiocellata-przewalskii* species complex resulting from partitioned Bayesian analysis based on barcoding *COI* haplotypes

Numbers beside nodes indicate Bayesian posterior probabilities (BPP). Colored symbols correspond to Figure 1 and those in Orlova et al. (2017), except light green branches and light green circle, which represent samples collected in this study.


Fig. S4



Supplementary Figure S4 Maximum-likelihood (ML) tree of *E. multiocellata-przewalskii* species complex resulting from partitioned ML analysis based on barcoding *COI* haplotypes

Numbers beside nodes indicate bootstrap support proportion (BSP). Colored symbols correspond to Figure 1 and those in Orlova et al. (2017), except light green branches and light green circle, which represent samples collected in this study.

Figure S5



Supplementary Figure S5 Automatic Barcode Gap Discovery (ABGD) species delimitation

A: Frequency histogram of Kimura 2-parameter (K2P) pairwise distances for all barcoding mitochondrial *COI* haplotypes. B: Partitions under different prior intraspecific divergences.

Supplementary Table S1 List of analyzed specimens, along with their sex, geographic origin and GenBank accession number

Taxon	Voucher No.	Sex	Population	Haplotype No.	Locality	Longitude	Latitude	GenBank No.	Reference
<i>E. dzungarica</i>	KZL91	♀	56	Hap 10	near the bank of Irtysh river, Bastaushy, Kokpekti District, Kazakhstan	83.42	48.82	MW172548	This study
<i>E. dzungarica</i>	KZL96	♀	56	Hap 10	near the bank of Irtysh river, Bastaushy, Kokpekti District, Kazakhstan	83.42	48.82	MW172549	This study
<i>E. dzungarica</i>	KZL97	♂	56	Hap 10	near the bank of Irtysh river, Bastaushy, Kokpekti District, Kazakhstan	83.42	48.82	MW172550	This study
<i>E. dzungarica</i>	KZL98	♂	56	Hap 15	near the bank of Irtysh river, Bastaushy, Kokpekti District, Kazakhstan	83.42	48.82	MW172551	This study
<i>E. dzungarica</i>	KZL99	♂	56	Hap 10	near the bank of Irtysh river, Bastaushy, Kokpekti District, Kazakhstan	83.42	48.82	MW172552	This study
<i>E. dzungarica</i>	KZL100	♀	56	Hap 10	near the bank of Irtysh river, Bastaushy, Kokpekti District, Kazakhstan	83.42	48.82	MW172542	This study
<i>E. dzungarica</i>	KZL111	♀	57	Hap 11	near the bank of Irtysh river, Zaisan desert, Kazakhstan	85.08	47.95	MW172543	This study
<i>E. dzungarica</i>	KZL113	♀	57	Hap 12	near the bank of Irtysh river, Zaisan desert, Kazakhstan	85.08	47.95	MW172544	This study
<i>E. dzungarica</i>	KZL114	♀	57	Hap 3	near the bank of Irtysh river, Zaisan desert, Kazakhstan	85.08	47.95	MW172545	This study
<i>E. dzungarica</i>	KZL115	♀	57	Hap 13	near the bank of Irtysh river, Zaisan desert, Kazakhstan	85.08	47.95	MW172546	This study
<i>E. dzungarica</i>	KZL116	♂	57	Hap 14	near the bank of Irtysh river, Zaisan desert, Kazakhstan	85.08	47.95	MW172547	This study

<i>E. dzungarica</i>	Guo8275	♀	58	Hap 3	Western Habahe County, Xinjiang, China	85.69	48.01	MW172538	This study
<i>E. dzungarica</i>	Guo8276	♂	58	Hap 5	Western Habahe County, Xinjiang, China	85.69	48.01	MW172539	This study
<i>E. dzungarica</i>	Guo8278	♀	58	Hap 5	Western Habahe County, Xinjiang, China	85.69	48.01	MW172540	This study
<i>E. dzungarica</i>	Guo8279	♂	58	Hap 9	Western Habahe County, Xinjiang, China	85.69	48.01	MW172541	This study
<i>E. dzungarica</i>	Guo3034	♂	59	Hap 3	Zaisan desert, Jeminay County, Xinjiang, China	85.60	47.71	MW172526	This study
<i>E. dzungarica</i>	Guo3035	♂	59	Hap 3	Zaisan desert, Jeminay County, Xinjiang, China	85.60	47.71	MW172527	This study
<i>E. dzungarica</i>	Guo3036	♂	59	Hap 4	Zaisan desert, Jeminay County, Xinjiang, China	85.60	47.71	MW172528	This study
<i>E. dzungarica</i>	Guo3023	♂	60	Hap 1	Southeast Jeminay County, Xinjiang, China	86.15	47.28	MW172523	This study
<i>E. dzungarica</i>	Guo3024	♀	60	Hap 1	Southeast Jeminay County, Xinjiang, China	86.15	47.28	MW172524	This study
<i>E. dzungarica</i>	Guo3026	♂	60	Hap 2	Southeast Jeminay County, Xinjiang, China	86.15	47.28	MW172525	This study
<i>E. dzungarica</i>	Guo8240	♂	61	Hap 8	Southeast Habahe County, Xinjiang, China	86.58	47.94	MW172534	This study
<i>E. dzungarica</i>	Guo8241	♂	61	Hap 5	Southeast Habahe County, Xinjiang, China	86.58	47.94	MW172535	This study
<i>E. dzungarica</i>	Guo8242	♀	61	Hap 5	Southeast Habahe County, Xinjiang, China	86.58	47.94	MW172536	This study
<i>E. dzungarica</i>	Guo8243	♀	61	Hap 5	Southeast Habahe County, Xinjiang, China	86.58	47.94	MW172537	This study
<i>E. dzungarica</i>	Guo3096	♀	62	Hap 5	Baishashan desert, Habahe County, Xinjiang, China	86.90	48.03	MW172529	This study
<i>E. dzungarica</i>	Guo3097	♂	62	Hap 6	Baishashan desert, Habahe County, Xinjiang, China	86.90	48.03	MW172530	This study
<i>E. dzungarica</i>	Guo3098	♀	62	Hap 7	Baishashan desert, Habahe County, Xinjiang, China	86.90	48.03	MW172531	This study
<i>E. dzungarica</i>	Guo3099	♂	62	Hap 5	Baishashan desert, Habahe County, Xinjiang, China	86.90	48.03	MW172532	This study

<i>E. dzungarica</i>	Guo3100	♀	62	Hap 5	Baishashan desert, Habahe County, Xinjiang, China	86.90	48.03	MW172533	This study
<i>E. szczerbaki</i>	ZMMU R-14342-1	\	1	Hap 46	Naryn Prov., Naryn Distr., N from Naryn, Kyrgyzstan	75.98	41.48	KY366618	Orlova et al. (2017)
<i>E. szczerbaki</i>	ZMMU R-14342-2	\	1	Hap 47	Naryn Prov., Naryn Distr., N from Naryn, Kyrgyzstan	75.98	41.48	KY366619	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-14341-2	\	2	Hap 44	Naryn Prov., Kochkor Distr., S from Kochkor, Kyrgyzstan	75.66	42.08	KY366616	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-14341-3	\	2	Hap 45	Naryn Prov., Kochkor Distr., S from Kochkor, Kyrgyzstan	75.66	42.08	KY366617	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-14339-1	\	3	Hap 41	Naryn Prov., Kochkor Distr., Kochkor, Kyrgyzstan	75.75	42.22	KY366615	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12556-1	\	4	Hap 37	Issyk-Kul Prov., env. of Balykchy, road to Akolen, Kyrgyzstan	76.17	42.35	KY366611	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12556-2	\	4	Hap 37	Issyk-Kul Prov., env. of Balykchy, road to Akolen, Kyrgyzstan	76.17	42.35	KY366612	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12557-1	\	4	Hap 43	Issyk-Kul Prov., env. of Balykchy, road to Akolen, Kyrgyzstan	76.17	42.35	KY366613	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12557-2	\	4	Hap 43	Issyk-Kul Prov., env. of Balykchy, road to Akolen, Kyrgyzstan	76.17	42.35	KY366614	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12427-1	\	5	Hap 42	Issyk-Kul Prov., NW bank of issyk-Kul lake, 20-25 km N from Toru Agyyr, Kungei-Alatau Mts., Kyrgyzstan	76.41	42.58	KY366609	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12427-2	\	5	Hap 42	Issyk-Kul Prov., NW bank of issyk-Kul lake, 20-25 km N from Toru Agyyr, Kungei-Alatau Mts., Kyrgyzstan	76.41	42.58	KY366610	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-14338-1	\	6	Hap 41	Issyk-Kul Prov., 100 km SW from Karakol,	77.18	42.16	KY366607	Orlova et al. (2017)

Kaji-Say env., Kyrgyzstan									
<i>E. stummeri</i>	ZMMU R-14338-2	\	6	Hap 41	Issyk-Kul Prov., 100 km SW from Karakol, Kaji-Say env., Kyrgyzstan	77.18	42.16	KY366608	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-14335-2	\	7	Hap 39	Issyk-Kul Prov., E bank of issyk-Kul lake, env. Of Karakol, Kyrgyzstan	78.36	42.46	KY366604	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-14335-3	\	7	Hap 40	Issyk-Kul Prov., E bank of issyk-Kul lake, env. Of Karakol, Kyrgyzstan	78.36	42.46	KY366605	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-14335-4	\	7	Hap 40	Issyk-Kul Prov., E bank of issyk-Kul lake, env. Of Karakol, Kyrgyzstan	78.36	42.46	KY366606	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-1a	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhstan	79.32	42.97	KY366589	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-2	\	8	Hap 35	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhstan	79.32	42.97	KY366590	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-2a	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen	79.32	42.97	KY366591	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-3	\	8	Hap 36	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhstan	79.32	42.97	KY366592	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-4	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhstan	79.32	42.97	KY366593	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-5	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhstan	79.32	42.97	KY366594	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-5a	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhstan	79.32	42.97	KY366595	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-6	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhstan	79.32	42.97	KY366596	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-7	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts.	79.32	42.97	KY366597	Orlova et al. (2017)

					Foothills, 7-8 km E from Kegen, Kazakhsan				
<i>E. stummeri</i>	ZMMU R-12551-8	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhsan	79.32	42.97	KY366598	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12551-10	\	8	Hap 34	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhsan	79.32	42.97	KY366599	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-1 2495-1	\	8	Hap 37	Almaty Prov., Rayimbek Distr., Ketmen Mts. Foothills, 7-8 km E from Kegen, Kazakhsan	79.32	42.97	KY366600	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12552-1	\	9	Hap 38	Almaty Prov., Rayimbek Distr., central Tian Shan Mts., 15 km S from Tuzkol Lake, Zhabyrtau Mt., Kazakhstan	80.08	42.92	KY366601	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12552-2	\	9	Hap 38	Almaty Prov., Rayimbek Distr., central Tian Shan Mts., 15 km S from Tuzkol Lake, Zhabyrtau Mt., Kazakhstan	80.08	42.92	KY366602	Orlova et al. (2017)
<i>E. stummeri</i>	ZMMU R-12552-4	\	9	Hap 38	Almaty Prov., Rayimbek Distr., central Tian Shan Mts., 15 km S from Tuzkol Lake, Zhabyrtau Mt., Kazakhstan	80.08	42.92	KY366603	Orlova et al. (2017)
<i>Eremias</i> sp. 1	ZMMU R-14329-1	\	10	Hap 50	Xinjiang Prov., 60 km NE from Aksu, China	80.79	41.54	KY366626	Orlova et al. (2017)
<i>Eremias</i> sp. 1	ZMMU R-14330-1	\	11	Hap 51	Xinjiang Prov., 75 km NE from Aksu, China	80.83	41.74	KY366627	Orlova et al. (2017)
<i>Eremias</i> sp. 1	ZMMU R-14330-2	\	11	Hap 52	Xinjiang Prov., 75 km NE from Aksu, China	80.83	41.74	KY366628	Orlova et al. (2017)
<i>Eremias</i> sp. 1	ZMMU R-14330-3	\	11	Hap 53	Xinjiang Prov., 75 km NE from Aksu, China	80.83	41.74	KY366629	Orlova et al. (2017)
<i>Eremias</i> sp. 1	ZMMU R-14327-1	\	12	Hap 49	Xinjiang Prov., 35 km NE from Aksu, China	81.05	41.40	KY366625	Orlova et al. (2017)
<i>Eremias</i> sp. 1	ZMMU R-14328-1	\	13	Hap 54	Xinjiang Prov., 89 km NE from Aksu, China	81.21	41.56	KY366630	Orlova et al. (2017)
<i>Eremias</i> sp. 1	ZMMU R-14328-2	\	13	Hap 55	Xinjiang Prov., 89 km NE from Aksu, China	81.21	41.56	KY366631	Orlova et al. (2017)
<i>Eremias</i> sp. 1	ZMMU R-14328-3	\	13	Hap 56	Xinjiang Prov., 89 km NE from Aksu, China	81.21	41.56	KY366632	Orlova et al. (2017)

<i>E. yarkandensis</i>	ZMMU R-14344-1	\	14	Hap 48	Osh Prov., vicinity of Nura, Kyrgyzstan	73.87	39.65	KY366620	Orlova et al. (2017)
<i>E. yarkandensis</i>	ZMMU R-14344-2	\	14	Hap 48	Osh Prov., vicinity of Nura, Kyrgyzstan	73.87	39.65	KY366621	Orlova et al. (2017)
<i>E. yarkandensis</i>	ZMMU R-14344-3	\	14	Hap 48	Osh Prov., vicinity of Nura, Kyrgyzstan	73.87	39.65	KY366622	Orlova et al. (2017)
<i>E. yarkandensis</i>	ZMMU R-14344-4	\	14	Hap 48	Osh Prov., vicinity of Nura, Kyrgyzstan	73.87	39.65	KY366623	Orlova et al. (2017)
<i>E. yarkandensis</i>	ZMMU R-14344-5	\	14	Hap 48	Osh Prov., vicinity of Nura, Kyrgyzstan	73.87	39.65	KY366624	Orlova et al. (2017)
<i>E. cf. buechneri</i>	ZMMU R-8910-1a	\	15	Hap 28	Xinjiang Prov., Qarqan (Chemo) Distr., Altintag Mt., Chinbulak, 60 km S from Turav, China	86.05	37.51	KY366574	Orlova et al. (2017)
<i>E. cf. buechneri</i>	ZMMU R-8910-1b	\	15	Hap 28	Xinjiang Prov., Qarqan (Chemo) Distr., Altintag Mt., Chinbulak, 60 km S from Turav, China	86.05	37.51	KY366575	Orlova et al. (2017)
<i>Eremias</i> sp. 2	\	\	16	Hap 26	Qinhai Prov., Delingha env., China	97.35	37.35	KY366572	Orlova et al. (2017)
<i>Eremias</i> sp. 2	\	\	17	Hap 27	Qinhai Prov., Gonghe (Qiabugia), Chekhou, env. Of Kyiking (Chin-Kon), China	100.75	36.20	KY366573	Orlova et al. (2017)
<i>E. cf. multiocellata</i>	\	\	18	Hap 65	Gansu Prov., env. of Shandan, China			KY366641	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-13209	\	19	Hap 17	Inner Mongolia Prov., 30 km W from Bayanhot	101.27	39.12	KY366551	Orlova et al. (2017)
<i>E. przewalskii</i>	\	\	20	Hap 16	Gansu Prov., Minqin County, Wuwei, China	103.11	38.63	KM507330	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-13214	\	21	Hap 18	Inner Mongolia Prov., Alashan desert, 150 km W from Bayanmod, China	105.59	40.41	KY366552	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-12046-1	\	22	Hap 19	Bayankhongor Aimaq, Zhinst, Mongolia	100.21	45.62	KY366553	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-13038-1	\	23	Hap 19	Govi-Altai Aimaq, Biger valley, Mongolia	97.4	45.73	KY366554	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-13038-3	\	23	Hap 19	Govi-Altai Aimaq, Biger valley, Mongolia	97.4	45.73	KY366555	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-13038-4	\	23	Hap 19	Govi-Altai Aimaq, Biger valley, Mongolia	97.4	45.73	KY366556	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-12842-1	\	24	Hap 19	Govi-Aitai Aimaq, 3 km W from Biger Sum, Mongolia	97.15	45.72	KY366557	Orlova et al. (2017)

<i>E. przewalskii</i>	ZMMU R-12842-2	\	24	Hap 19	Govi-Aitai Aimaq, 3 km W from Biger Sum, Mongolia	97.15	45.72	KY366558	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-12888-1	\	25	Hap 20	Govi-Aitai Aimaq, Tsogt, Dzahuin-Govi, 10 km W Bayan-Toro, Mongolia	96.65	45.67	KY366559	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-12888-2	\	25	Hap 20	Govi-Aitai Aimaq, Tsogt, Dzahuin-Govi, 10 km W Bayan-Toro, Mongolia	96.65	45.67	KY366560	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-13060-1	\	26	Hap 19	Tuva Republic, Ovyurskiy Distr., Tere-Khollake, Tsuger-Els sands, Russia	93.21	50.68	KY366561	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-13060-2	\	26	Hap 19	Tuva Republic, Ovyurskiy Distr., Tere-Khollake, Tsuger-Els sands, Russia	93.21	50.68	KY366562	Orlova et al. (2017)
<i>E. przewalskii</i>	ZMMU R-13060-3	\	26	Hap 19	Tuva Republic, Ovyurskiy Distr., Tere-Khollake, Tsuger-Els sands, Russia	93.21	50.68	KY366563	Orlova et al. (2017)
<i>E. przewalskii</i>	ENS-T07-1	\	26	Hap 19	Tuva Republic, Ovyurskiy Distr., Tere-Khollake, Tsuger-Els sands, Russia	93.21	50.68	KY366564	Orlova et al. (2017)
<i>E. przewalskii</i>	ENS-T07-2	\	26	Hap 19	Tuva Republic, Ovyurskiy Distr., Tere-Khollake, Tsuger-Els sands, Russia	93.21	50.68	KY366565	Orlova et al. (2017)
<i>E. cf. reticulata</i>	ZMMU R-12855-1	\	27	Hap 22	Gobi-Altai Aimaq, 4 km NW from Altai Sum, env. Danshig-Khuduk, Mongolia	95.86	44.64	KY366567	Orlova et al. (2017)
<i>E. cf. reticulata</i>	ZMMU R-12855-2	\	27	Hap 23	Gobi-Altai Aimaq, 4 km NW from Altai Sum, env. Danshig-Khuduk, Mongolia	95.86	44.64	KY366568	Orlova et al. (2017)
<i>E. cf. reticulata</i>	ZMMU R-12856-1	\	27	Hap 23	Gobi-Altai Aimaq, 4 km NW from Altai Sum, env. Danshig-Khuduk, Mongolia	95.86	44.64	KY366569	Orlova et al. (2017)
<i>E. cf. reticulata</i>	MNUE-Em-S	\	28	Hap 21	Bayankhongor Aimaq, Tsagan Bogdo Uul (Mt.)	98.82	42.88	KY366566	Orlova et al. (2017)
<i>E. cf. przewalskii</i>	ZMMU R-12858-1	\	29	Hap 24	ömnögovi Aimaq, 8-10 km S from Bayandalai Sum, Mongolia	103.45	43.39	KY366570	Orlova et al. (2017)
<i>E. cf. przewalskii</i>	ZMMU R-12859-1	\	29	Hap 25	ömnögovi Aimaq, 8-10 km S from Bayandalai	103.45	43.39	KY366571	Orlova et al. (2017)

Sum, Mongolia										
<i>Eremias</i> sp. 3	\	\	30	Hap 64	Gansu Prov., env. of Lanzhou, Baiyin, China	104.21	36.46	KJ664798	Li and Song (unpublished data)	
<i>Eremias</i> sp. 3	\	\	31	Hap 63	Gansu Prov., Minqin County, China	103.11	38.63	NC025304	Tong et al. (2014)	
<i>Eremias</i> sp. 3	ZMMU R-13208	\	32	Hap 62	Inner Mongolia Prov., 80 km SW Sayan Nur, China	103.87	40.00	KY366640	Orlova et al. (2017)	
<i>Eremias</i> sp. 3	ZMMU R-13207	\	33	Hap 60	Inner Mongolia Prov., 120 km W from Wuhai, China	105.45	39.53	KY366638	Orlova et al. (2017)	
<i>Eremias</i> sp. 3	ZMMUR-13206	\	34	Hap 61	Inner Mongolia Prov., Alashan desert, 40 km S from Bayan-Hot, China	105.62	38.67	KY366639	Orlova et al. (2017)	
<i>Eremias</i> sp. 3	ZMMU R-13205	\	35	Hap 59	Inner Mongolia Prov., W Ordos, 30 km SW from Bayan-Us, China	108.50	40.02	KY366637	Orlova et al. (2017)	
<i>Eremias</i> sp. 3	ZMMU R-13215	\	36	Hap 58	Inner Mongolia Prov., 50 km S from Baotou, China	110.00	40.28	KY366636	Orlova et al. (2017)	
<i>Eremias</i> sp. 3	ZMMU R-12610-1a	\	37	Hap 57	Dornogovi Aimaq, Sainshand Sum, vicinity of Saishand, Mongolia	110.10	44.89	KY366633	Orlova et al. (2017)	
<i>Eremias</i> sp. 3	ZMMU R-12610-1b	\	37	Hap 57	Dornogovi Aimaq, Sainshand Sum, vicinity of Saishand, Mongolia	110.10	44.89	KY366634	Orlova et al. (2017)	
<i>Eremias</i> sp. 3	ZMMU R-14523	\	38	Hap 57	Dornogovi Aimaq, Sainshand Sum, ca. 100 km NE from örgön, Mongolia	111.79	45.18	KY366635	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-12863-1	\	39	Hap 72	Umnugovi Aimaq, 102 km NW from Dalandzadagat, Tsogt-Ovu, Mongolia	105.32	44.42	KY366651	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-12863-2	\	39	Hap 73	Umnugovi Aimaq, 102 km NW from Dalandzadagat, Tsogt-Ovu, Mongolia	105.32	44.42	KY366652	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-12952a	\	40	Hap 70	Dundgovi Aimaq, env. of Deren, Mongolia	106.54	46.59	KY366647	Orlova et al. (2017)	

<i>E. multiocellata</i>	ZMMU R-12952b	\	40	Hap 70	Dundgovi Aimaq, env. of Deren, Mongolia	106.54	46.59	KY366648	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12608-1	\	41	Hap 71	Dundgovi Aimaq, env. of ölziit Sum, Mongolia	102.51	48.03	KY366649	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12608-2	\	41	Hap 71	Dundgovi Aimaq, env. of ölziit Sum, Mongolia	102.51	48.03	KY366650	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12045-1	\	42	Hap 74	Bayankhongor Aimaq, Zhinst, Mongolia	100.21	45.62	KY366653	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12045-2	\	42	Hap 74	Bayankhongor Aimaq, Zhinst, Mongolia	100.21	45.62	KY366654	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12841-1	\	43	Hap 74	Bayankhongor Aimaq, N bank of Bööntsagan Nuur Lake, Mongolia	99.12	45.67	KY366655	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12841-2	\	43	Hap 74	Bayankhongor Aimaq, N bank of Bööntsagan Nuur Lake, Mongolia	99.12	45.67	KY366656	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-13039-1	\	44	Hap 74	Govi-Altai Aimaq, Delger, Mongolia	98.05	45.93	KY366657	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12843-1	\	45	Hap 75	Govi-Altai Aimaq, Shargyn-Govi, 2 km SW from Khalium Sum, Mongolia	96.12	45.93	KY366658	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12843-2	\	45	Hap 75	Govi-Altai Aimaq, Shargyn-Govi, 2 km SW from Khalium Sum, Mongolia	96.12	45.93	KY366659	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12081-1	\	46	Hap 76	Govi-Aitai Aimaq, Shargyn-Govi, 30 km S from Sharga Sum, Mongolia	95.23	46.36	KY366660	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-13058-1	\	47	Hap 66	Tuva Republic, Erzin Distr., Ubsu-Nw valley, 12 km SE from Yamalyg Mt., Russia	94.79	50.16	KY366642	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-13058-2	\	47	Hap 67	Tuva Republic, Erzin Distr., Ubsu-Nw valley, 12 km SE from Yamalyg Mt., Russia	94.79	50.16	KY366643	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-13058-3	\	47	Hap 68	Tuva Republic, Erzin Distr., Ubsu-Nw valley, 12 km SE from Yamalyg Mt., Russia	94.79	50.16	KY366644	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-13058-4	\	47	Hap 69	Tuva Republic, Erzin Distr., Ubsu-Nw valley, 12 km SE from Yamalyg Mt., Russia	94.79	50.16	KY366645	Orlova et al. (2017)

<i>E. multiocellata</i>	ZMMU R-13059-1	\	47	Hap 67	Tuva Republic, Erzin Distr., Ubsu-Nw valley, km SE from Yamalyg Mt., Russia	12	94.79	50.16	KY366646	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12844-1	\	48	Hap 74	Govi-Aitai Aimaq, Junggar Govi, 4 km W from Bidzh, Mongolia		93.58	45.59	KY366661	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12844-2	\	48	Hap 74	Govi-Aitai Aimaq, Junggar Govi, 4 km W from Bidzh, Mongolia		93.58	45.59	KY366662	Orlova et al. (2017)
<i>E. multiocellata</i>	ZMMU R-12853-1	\	49	Hap 74	Govi-Aitai Aimaq, Junggar Govi, Bugat, Mongolia	93.53	45.22	KY366663	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-12853-2	\	49	Hap 74	Govi-Aitai Aimaq, Junggar Govi, Bugat, Mongolia	93.53	45.22	KY366664	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-12848-1	\	50	Hap 74	Khovd Aimaq, 58 km SE from Altai Sum, E foothills of Sertengiyn-Khuvch-Ula, Mongolia	92.98	45.65	KY366665	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-12849-1	\	50	Hap 77	Khovd Aimaq, 58 km SE from Altai Sum, E foothills of Sertengiyn-Khuvch-Ula, Mongolia	92.98	45.65	KY366666	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-12849-2	\	50	Hap 74	Khovd Aimaq, 58 km SE from Altai Sum, E foothills of Sertengiyn-Khuvch-Ula, Mongolia	92.98	45.65	KY366667	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-12847-1	\	51	Hap 75	Khovd Aimaq, I km SW from Altai Sum, Mongolia	92.25	45.80	KY366668	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-13132-1	\	52	Hap 74	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366669	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-13132-2	\	52	Hap 74	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366670	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-13132-3	\	52	Hap 74	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366671	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-13132-4	\	52	Hap 74	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366672	Orlova et al. (2017)	
<i>E. multiocellata</i>	ZMMU R-13132-5	\	52	Hap 74	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366673	Orlova et al. (2017)	

<i>E. dzungarica</i>	ZMMU R-12845-2	\	52	Hap 31	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366580	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12845-4	\	52	Hap 31	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366581	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12845-5	\	52	Hap 32	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366582	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12845-6	\	52	Hap 31	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366583	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12845-7	\	52	Hap 31	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366584	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12845-8	\	52	Hap 31	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366585	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12845-9	\	52	Hap 31	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366586	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12845-10	\	52	Hap 31	Khovd Aimaq, 7 km SW from Uyench Sum, Mongolia	93.61	46.93	KY366587	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12862-1	\	53	Hap 31	Khovd Aimaq, Bulgan Sum, Bayan-Mod, 11 km W from Ikher-Toli, Mongolia	92.91	47.06	KY366578	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12862-3	\	53	Hap 31	Khovd Aimaq, Bulgan Sum, Bayan-Mod, 11 km W from Ikher-Toli, Mongolia	92.91	47.06	KY366579	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-12550-1	\	54	Hap 33	Khovd Aimaq, 24 km N from Uyench Sum, Mongolia	92.05	46.27	KY366588	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-11989-1	\	55	Hap 29	East-Kazakhstan Prov., Aigyrkum sands, 5-7 km SW from Buran	84.89	47.98	KY366576	Orlova et al. (2017)
<i>E. dzungarica</i>	ZMMU R-11989-2	\	55	Hap 30	East-Kazakhstan Prov., Aigyrkum sands, 5-7 km SW from Buran	84.89	47.98	KY366577	Orlova et al. (2017)

<i>E. argus barbouri</i>	ZMMU R-12605-1	\	\	\	\	\	\	\	KY366548	Orlova et al. (2017)
<i>E. brenchleyi</i>	\	\	\	\	Qianshan, Suzhou, Anhui Province, China	\	\	\	EF490071	Rui et al. (2009)
<i>E. nikolskii</i>	ZMMU R-11673-1	\	\	\	\	\	\	\	KY366550	Orlova et al. (2017)
<i>E. vermiculata</i>	ZMMU-R12047-1	\	\	\	\	\	\	\	KY366549	Orlova et al. (2017)

References

- Orlova VF, Poyarkov Jr NA, Chirikova MA, Nazarov RA, Munkhbaatar M, Munkhbayar K, Terbish K. 2017. MtDNA differentiation and taxonomy of Central Asian racerunners of *Eremias multiocellata*-*E. przewalskii* species complex (Squamata, Lacertidae). *Zootaxa*, **4282**(1): 1–42.
- Rui J, Wang Y, Nie L. 2009. The complete mitochondrial DNA genome of *Eremias brenchleyi* (Reptilia: Lacertidae) and its phylogeny position within squamata reptiles. *Amphibia-Reptilia*, **30**: 25–35

Supplementary Table S2 Kimura 2-parameter (K2P) distance summary for sequence divergence between barcoding mitochondrial *COI* sequences at species and genus level

	Individuals	Taxa	Comparisons	Min%	Mean%	Max%	SE%
Within species	155	12	2018	0.00	1.16	6.18	0.00
Within genus	156	1	10072	0.77	9.09	13.19	0.00

Supplementary Table S3 Kimura 2-parameter (K2P) distance summary of maximum intraspecific variation and minimum genetic distance to the nearest-neighbor species calculated from barcoding mitochondrial *COI* sequences for each species in *Eremias multiocellata-przewalskii* species complex

Species	Max Intra-Sp (%)	Nearest Species	Distance to NN (%)
<i>Eremias</i> sp. 2	0.31	<i>Eremias</i> . cf. <i>reticulata</i>	0.77
<i>Eremias</i> . cf. <i>reticulata</i>	0.31	<i>Eremias</i> sp. 2	0.77
<i>Eremias przewalskii</i>	2.19	<i>Eremias</i> . cf. <i>przewalskii</i>	1.08
<i>Eremias</i> . cf. <i>przewalskii</i>	0.31	<i>Eremias przewalskii</i>	1.08
<i>Eremias</i> cf. <i>buechneri</i>	0	<i>Eremias yarkandensis</i>	1.48
<i>Eremias yarkandensis</i>	0	<i>Eremias</i> cf. <i>buechneri</i>	1.48
<i>Eremias</i> . <i>dzungarica</i>	2.99	<i>Eremias</i> cf. <i>buechneri</i>	1.87
<i>Eremias</i> sp. 1	2.14	<i>Eremias yarkandensis</i>	3.35
<i>Eremias multiocellata</i>	1.71	<i>Eremias</i> sp. 3	5.65
<i>Eremias</i> sp. 3	6.18	<i>Eremias multiocellata</i>	5.65
<i>Eremias</i> cf. <i>multiocellata</i>	N/A	<i>Eremias multiocellata</i>	6.85
<i>Eremias stummeri</i>	2.47	<i>Eremias szczerbaki</i>	7.58
<i>Eremias szczerbaki</i>	0.33	<i>Eremias stummeri</i>	7.58

Supplementary Table S4 Quantitative characteristics of specimens of *Eremias dzungarica* from this study and Orlova et al. (2017)

Values are given as ranges and means \pm SE. Morphological characters detected with significant sexual dimorphism are given separately for females and males.

Table S4.1

Metric and meristic traits	Females of <i>E. dzungarica</i> in this study			Females of <i>E. dzungarica</i> from Orlova et al. (2017)		
	N	Range	Mean \pm SE	N	Range	Mean \pm SE
SVL	15	46.55-64.46	57.79 \pm 1.443	2	61.40-64.50	\
TL	7	63.65-76.19	69.96 \pm 1.794	0	\	\
Ga	15	21.48-33.07	28.72 \pm 0.941	2	33.80-35.00	\
HL	15	11.59-15.40	13.64 \pm 0.279	2	14.00-15.00	\
HW	15	6.99-9.89	8.66 \pm 0.196	2	7.00-8.20	\
HH	15	5.47-8.07	6.96 \pm 0.215	2	6.00-7.20	\
NL	15	15.98-20.96	18.40 \pm 0.353	2	19.00-20.50	\
Pa	15	16.14-20.48	18.68 \pm 0.346	2	18.00-19.00	\
Pp	15	23.95-29.66	26.98 \pm 0.459	2	25.40-26.20	\
Dist.P.f						
m	15	3.52-5.42	4.73 \pm 0.117	2	1.60-1.70	\
Sq.c.cd	15	24-27	25.13 \pm 0.274	2	23-23	
Lpil.	15	10.74-14.15	12.34 \pm 0.236	0	\	\

Table S4.2

Metric and meristic traits	Males of <i>E. dzungarica</i> in this study			Males of <i>E. dzungarica</i> from Orlova et al. (2017)		
	N	Range	Mean \pm SE	N	Range	Mean \pm SE
SVL	15	56.14-75.72	63.44 \pm 1.378	5	52.00-65.00	58.72 \pm 2.506
TL	11	72.74-94.82	80.79 \pm 2.121	2	67.30-85.00	\
Ga	15	25.23-35.61	31.08 \pm 0.853	5	24.00-32.00	28.240 \pm 1.574
HL	15	15.10-18.98	16.45 \pm 0.343	5	13.60-18.00	15.88 \pm 0.779
HW	15	8.82-12.20	10.50 \pm 0.232	5	7.30-10.30	9.00 \pm 0.640
HH	15	7.16-10.72	8.84 \pm 0.272	5	5.00-8.30	6.86 \pm 0.616
NL	15	19.53-26.14	21.93 \pm 0.473	5	12.50-21.40	18.96 \pm 1.688
Pa	15	18.14-24.33	20.81 \pm 0.414	5	17.50-20.00	19.20 \pm 0.464
Pp	15	26.64-35.41	31.16 \pm 0.654	5	25.70-30.00	28.48 \pm 0.813
Dist.P.fm	15	3.52-6.34	4.68 \pm 0.219	5	1.70-2.50	2.06 \pm 0.129
Sq.c.cd	15	25-29	26.67 \pm 0.319	5	24-27	25.20 \pm 0.583
Lpil.	15	13.26-16.48	14.35 \pm 0.270	0	\	\

Table S4.3

Meristic traits	All specimens of <i>E. dzungarica</i> in this study			All specimens of <i>E. dzungarica</i> from Orlova et al. (2017)		
	N	Range	Mean ± SE	N	Range	Mean ± SE
Lab.total.L	30	9–12	10.83 ± 0.145	11	9–11	10.36 ± 0.203
Lab.total.R	30	9–12	10.60 ± 0.141	11	9–11	10.00 ± 0.234
Infralab.L	30	6–7	6.63 ± 0.089	12	6–8	6.83 ± 0.167
Infralab.R	30	6–8	6.77 ± 0.092	12	6v9	6.92 ± 0.260
Sq.	30	44–55	47.67 ± 0.456	12	46–52	48.42 ± 0.609
Ventr.	30	30–34	31.93 ± 0.185	12	30v32	30.92 ± 0.193
G.	30	20–26	22.93 ± 0.335	12	21–26	22.83 ± 0.474
Coll.	30	9–14	11.27 ± 0.267	11	7–13	11.00 ± 0.632
Lam.subdig.	30	22–25	23.53 ± 0.224	12	19–24	21.58 ± 0.379
P.fm.L	30	10–13	11.17 ± 0.167	12	9–15	12.17 ± 0.405
P.fm.R	30	9–14	11.33 ± 0.188	12	9–14	12.00 ± 0.369
Ventr.long.	30	14–18	16.30 ± 0.199	1	16	\
supracil.L	30	3–8	5.43 ± 0.196	0	\	\
supracil.R	30	4–7	5.47 ± 0.124	0	\	\
scal.f.p	30	7–12	9.70 ± 0.204	0	\	\

Supplementary Table S5 Morphological characters used for morphometric analysis and their corresponding abbreviations

Abbreviations	Description of morphological trait
SVL	Snout to vent length
TL	Tail length
Ga	Gleno-acetabular distance (from axilla to groin)
HL	Head length
HW	Head width
HH	Head height
Lpil.	Pileus Length (from rostrum to the posterior border of parietals)
NL	head length from snout tip to the anterior edge of collar (measured from ventral side)
Pa	Forelimb Length
Pp	Hindlimb Length
Dist.P.fm.	Distance between femoral pore rows
Lab.total.L	Number of supralabials on the left side
Lab.total.R	Number of supralabials on the right side
Infralab.L	Number of infralabials on the left side
Infralab.R	Number of infralabials on the right side
Sq.	Number of scales around middorsal
Ventr.	Number of transverse rows of ventral scales
G.	Number of gular scales along mid-line of throat
Coll.	Number of collar scales
Lam.subdig.	Number of subdigital lamellae on the 4th toe of hindlimb
P.fm.L	Number of femoral pores on the left side
P.fm.R	Number of femoral pores on the right side
Ventr.long.	Maximal number of longitudinal rows of ventral scales
Supracil.L	Number of supraciliary scales on the left side
Supracil.R	Number of supraciliary scales on the right side
Scal.f.p	Number of scales between femoral pore rows
Sq.c.cd.	Number of scales around the 9–10th tail ring